

PRESCRIBING RICCI CURVATURE ON COMPLEXIFIED SYMMETRIC SPACES

ROGER BIELAWSKI

The aim of this note is to prove

Theorem 1. *Let G/K be an irreducible symmetric space of compact type with K connected and G acting effectively on G/K , and let ρ be a real exact G -invariant $(1, 1)$ -form on the complexification $G^{\mathbb{C}}/K^{\mathbb{C}}$. Then there exists a G -invariant Kähler metric on $G^{\mathbb{C}}/K^{\mathbb{C}}$ whose Ricci form is ρ .*

Remarks: 1. The assumptions on G and K are fulfilled for simply connected symmetric spaces. In general, these assumptions guarantee that G is connected ([6], Thm. V.4.1), that G/K is globally symmetric ([6], Ex. VII.10), and that $K = \text{Ad}_G(K)$ (as $\text{Ad}_G(K) = K/K \cap Z(G)$).

2. The Kähler form obtained in Theorem 1 is exact.

The above result has been proved in [9] for symmetric spaces of rank 1 and in [2] for compact groups, i.e. for the case when $G = K \times K$ and K acts diagonally. For hermitian symmetric spaces and $\rho = 0$, Theorem 1 has also been known [4].

The proof given here is quite different from that given for group manifolds in [2]. We show that the complex Monge-Ampère equation on $G^{\mathbb{C}}/K^{\mathbb{C}}$ reduces, for G -invariant functions, to a real Monge-Ampère equation on the dual symmetric space G^*/K . We also show that the Monge-Ampère operator on non-compact symmetric spaces has a radial part, i.e. it is equal, for K -invariant functions, to another Monge-Ampère operator on the maximal abelian subspace of G^*/K . These facts, together with the theorem on K -invariant real Monge-Ampère equations proved in [3], yield Theorem 1.

1. RIEMANNIAN SYMMETRIC SPACES OF NON-COMPACT TYPE

Here we recall some facts about the geometry of Riemannian symmetric spaces. The standard reference for this section is [6].

Let G/K be a symmetric space of compact type with G acting effectively and K connected, and let G^*/K be its dual². If $\mathfrak{g}$, $\mathfrak{g}^*$ and $\mathfrak{k}$ denote the Lie algebras of G , G^* and K , then $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, $\mathfrak{g}^* = \mathfrak{k} \oplus i\mathfrak{p}$, where $[\mathfrak{k}, \mathfrak{p}] \subset \mathfrak{p}$ and $[\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}$. The restriction of the Killing form to $i\mathfrak{p}$ is positive definite and induces the Riemannian metric of G^*/K . Moreover, the Riemannian exponential mapping provides a diffeomorphism between $\mathfrak{p}$ and G^*/K . This can be viewed as the map:

$$p \mapsto e^{ip}K, \tag{1.1}$$

¹The complexification of a compact connected Lie group G is the connected group $G^{\mathbb{C}}$ whose Lie algebra is the complexification of the Lie algebra of G and which satisfies $\pi_1(G^{\mathbb{C}}) = \pi_1(G)$.

²Since K is connected and $K = \text{Ad}_G(K)$ (Remark 1), there exists a G^* such that (G^*, K) is a symmetric pair corresponding to $(\mathfrak{g}^*, \mathfrak{k})$.

where $p \in \mathfrak{p}$ and e is the group-theoretic exponential map for G^* . Thus we have two K -invariant Riemannian metrics on $\mathfrak{p} \simeq \mathbb{R}^n$: the Euclidean one given by the Killing form, and the negatively curved one given by the diffeomorphism (1.1).

Let $\mathfrak{a}$ be a maximal abelian subspace of $\mathfrak{p}$ and $\mathfrak{l}$ its centraliser in $\mathfrak{k}$. Let Σ the set of restricted roots and Σ^+ the set of restricted positive roots. For each $\alpha \in \Sigma$, let $\mathfrak{p}_\alpha$ (resp. $\mathfrak{k}_\alpha$) denote the subspace of $\mathfrak{p}$ (resp. of $\mathfrak{k}$) where each $(\text{ad } H)^2$, $H \in \mathfrak{a}$, acts with eigenvalue $\alpha(H)^2$. We have the direct decompositions

$$\mathfrak{p} = \mathfrak{a} + \sum_{\alpha \in \Sigma^+} \mathfrak{p}_\alpha, \quad \mathfrak{k} = \mathfrak{l} + \sum_{\alpha \in \Sigma^+} \mathfrak{k}_\alpha. \quad (1.2)$$

Let $\mathfrak{a}^+$ be an open Weyl chamber and let $\mathfrak{p}'$ be the union of K -orbits of points in $\mathfrak{a}^+$. Any K orbit in $\mathfrak{p}'$ is isomorphic to K/L where the Lie algebra of L is $\mathfrak{l}$. Moreover, we have the diffeomorphism:

$$\mathfrak{a}^+ \times K/L \rightarrow \mathfrak{p}', \quad (h, k) \mapsto \text{Ad}(k)h. \quad (1.3)$$

We now wish to write the two K -invariant metrics on $\mathfrak{p}$ in coordinates given by this diffeomorphism. Let $\sum dr_i^2$ be the Killing metric on $\mathfrak{a}^+$ (the r_i can be viewed as K -invariant functions on $\mathfrak{p}'$). For each $\mathfrak{k}_\alpha$, choose a basis $X_{\alpha, m}$ (m runs from 1 to twice the multiplicity of α) of vectors orthonormal for the Killing form and denote by $\theta_{\alpha, m}$ the corresponding basis of invariant 1-forms on K/L . We have

Proposition 1.1. *Let g_0 be the Euclidean metric on $\mathfrak{p}$, given by the restriction of the Killing form, and let g be the negatively curved symmetric metric on $\mathfrak{p}$ given by the diffeomorphism (1.1). Then, under the diffeomorphism (1.3) the metrics g_0 and g can be written in the form*

$$\sum_i dr_i^2 + \sum_{(\alpha, m)} F(\alpha(r)) \theta_{(\alpha, m)}^2, \quad (1.4)$$

where $F(z) = z^2$ for g_0 , and $F(z) = \sinh^2(z)$ for g .

Proof. Since all these metrics are K -invariant, it is enough to compute them at points of $\mathfrak{a}^+$. Let H be such a point and let (h, ρ) , $h \in \mathfrak{a}$, $\rho \in T_{[1]}K/L$, be a tangent vector to $\mathfrak{a}^+ \times K/L$ at $(H, [1])$. The vector ρ can be identified with an element of $\sum \mathfrak{k}_\alpha \subset \mathfrak{k}$. The corresponding (under (1.3)) tangent vector at $H \in \mathfrak{p}'$ is $h + [\rho, H]$. Computing the Killing form of this vector yields the formula (1.4) with $F(z) = z^2$ for g_0 . The formula for g follows from a similar computation, using the expression for the differential of the map (1.1) given in [6], Theorem IV.4.1. $\square$

2. MONGE-AMPÈRE EQUATION ON SYMMETRIC SPACES

Let (M, g) be a Riemannian manifold and $u : M \rightarrow \mathbb{R}$ a smooth function. Then the Hessian of u is the symmetric $(0, 2)$ -tensor Ddu where D is the Levi-Civita connection of g . In local coordinates x_i , Ddu is represented by the matrix

$$H_{ij} = \frac{\partial^2 u}{\partial x_i \partial x_j} - \sum_k \Gamma_{ij}^k \frac{\partial u}{\partial x_k}. \quad (2.1)$$

We say that the function u is g -convex (resp. strictly g -convex), if Ddu is non-negative (resp. positive) definite. The Monge-Ampère equation on the manifold (M, g) is then

$$\mathbf{M}_g(u) := (\det g)^{-1} \det Ddu = f \quad (2.2)$$

where f is a given function.

Let $(G^*/K, g)$ be a symmetric space of non-compact type given by a Cartan decomposition $\mathfrak{g}^* = \mathfrak{k} + i\mathfrak{p}$. As in the previous section, we identify $M = G^*/K$ with $\mathfrak{p}$ and denote by g_0 the (flat) metric given by restricting the Killing form to $\mathfrak{p}$. We have:

Theorem 2.1. *Let $M \simeq \mathfrak{p}$ be a symmetric space of noncompact type and let u be a K -invariant (smooth) function on M . Then*

- (1) *u is g -convex if and only if u is g_0 -convex (i.e. convex in the usual sense on $\mathfrak{p}$).*
- (2) *The following equality of Monge-Ampère operators holds:*

$$\mathbf{M}_g(u) = F \cdot \mathbf{M}_{g_0}(u),$$

where $F : M \rightarrow \mathbb{R}$ is a positive K -invariant smooth function depending only on M .

We have proved in [3] a theorem on the existence and regularity of K -invariant solutions to Monge-Ampère equations on $\mathbb{R}^n$. From this we immediately obtain

Corollary 2.2. *Let $(G^*/K, g)$ be an irreducible symmetric space of noncompact type and let f be a positive smooth K -invariant function on G^*/K . Then the Monge-Ampère equation (2.2) has a global smooth K -invariant strictly g -convex solution. $\square$*

We shall now prove Theorem 2.1. In fact we shall prove it in the following, more general situation. Suppose that we are given a K -invariant metric on $\mathfrak{p}$ whose pullback under (1.3) can be written as (cf. (1.4)):

$$\sum_i dr_i^2 + \sum_{(\alpha, m)} F_{(\alpha, m)}(\alpha(r)) \theta_{(\alpha, m)}^2, \quad (2.3)$$

where $F_{(\alpha, m)} : \mathbb{R} \rightarrow \mathbb{R}$ are smooth functions vanishing at the origin such that $z^{-1} \frac{dF_{(\alpha, m)}}{dz}$ is smooth and positive everywhere. Proposition 1.1 implies that the symmetric metric on G^*/K is of this form. We claim that Theorem 2.1 holds for any metric g of the form (2.3).

In order to simplify the notation, let us write j for the index (α, m) and α_j for α if $j = (\alpha, m)$. The metric g can be now written as

$$\sum_i dr_i^2 + \sum_j F_j(\alpha_j(r)) \theta_j^2.$$

We recall the following formula:

$$2Ddu = L_{\nabla u} g,$$

where L is the Lie derivative and ∇u is the gradient of u with respect to the metric g . On the other hand, for any $(0, 2)$ -tensor g and vector fields X, Y, Z , we have:

$$(L_X g)(Y, Z) = X.g(Y, Z) - g([X, Y], Z) - g(Y, [X, Z]).$$

We now compute $L_{\nabla u} g$ on $\mathfrak{p}'$ with respect to the basis vector fields $\partial/\partial r_i$, X_j , where X_j are dual to θ_j . Here u is a K -invariant function. The gradient of u is just $\sum \frac{\partial u}{\partial r_i} \frac{\partial}{\partial r_i}$, in particular it is independent of the functions F_j . It follows

immediately that $(L_{\nabla u} g)(\partial/\partial r_i, X_j) = 0$ and that the matrix $(L_{\nabla u} g)(X_j, X_k)$ is equal to $\nabla u \cdot g(X_j, X_k)$ and hence it is diagonal with the (jj) -entry equal to

$$\nabla u(F_j(\alpha_j(r))) = \frac{dF_j}{dz} \Big|_{z=\alpha_j(r)} \alpha_j(\nabla_0 \bar{u}).$$

Here $\nabla_0 \bar{u} = \sum \frac{\partial u}{\partial r_i} \frac{\partial}{\partial r_i}$ is the gradient of u restricted to the Euclidean space $\mathfrak{a} = \mathbb{R}^n$ in coordinates r_i , and viewed as a map from $\mathbb{R}^n$ to itself.

Theorem 2.1 with the more general metric (2.3) follows easily with the function F given explicitly by

$$F = \frac{\prod \alpha_j(r)}{\prod F_j(\alpha_j(r))} \prod \left(\frac{1}{2} \frac{dF_j}{dz} \right)_{z=\alpha_j(r)}.$$

Observe that the assumptions on the F_j guarantee that F extends to a smooth positive function on $\mathfrak{p}$.

3. PROOF OF THE MAIN THEOREM

Let G/K be a locally symmetric space of compact type with K connected. There is a canonical isomorphism between $G^{\mathbb{C}}/K^{\mathbb{C}}$ and $G \times_K \mathfrak{p}$ (i.e. the tangent bundle of G/K) given by the map:

$$G \times \mathfrak{p} \rightarrow G^{\mathbb{C}} \rightarrow G^{\mathbb{C}}/K^{\mathbb{C}}, \quad (g, p) \mapsto ge^{ip}. \quad (3.1)$$

This isomorphism can be viewed in many ways: as an example of Mostow fibration [8], as given via Kähler reduction of $G^{\mathbb{C}} \simeq G \times \mathfrak{g}$ by the group K [5], or as given by the adapted complex structure construction [7] which provides a canonical diffeomorphism between the tangent bundle of G/K and a complexification of G/K . In any case it provides a fibration

$$\pi : G^{\mathbb{C}}/K^{\mathbb{C}} \rightarrow G/K. \quad (3.2)$$

The fibers of this projection can be identified with $\mathfrak{p}$ via the map (3.1). In particular, the fiber over $[1]$ is given by the $K^{\mathbb{C}}$ -orbits of elements $e^{ip}, p \in \mathfrak{p}$. We shall relate G -invariant plurisubharmonic functions on $G^{\mathbb{C}}/K^{\mathbb{C}}$ to convex functions on this fiber (see [1] for a different approach to this).

For a function w on a complex manifold one defines its Levy form Lw to be the Hermitian $(0, 2)$ tensor given in local coordinates as

$$\frac{\partial^2 w}{\partial z_k \partial \bar{z}_l} dz_k \otimes d\bar{z}_l. \quad (3.3)$$

This form does not depend on the choice of local coordinates. We shall compute this form for a G -invariant function w on $G^{\mathbb{C}}/K^{\mathbb{C}}$. It is enough to compute it at points $e^{ip}, p \in \mathfrak{p}$. First of all, we choose local holomorphic coordinates at such a point:

Lemma 3.1. *In a neighbourhood of a point $e^{ip}, p \in \mathfrak{p}$, complex coordinates are provided by the map $\mathfrak{p}^{\mathbb{C}} \rightarrow G^{\mathbb{C}} \rightarrow G^{\mathbb{C}}/K^{\mathbb{C}}, (a + ib) \mapsto e^{a+ib}e^{ip}$.*

Proof. We have to show that e^{a+ib} does not belong to the isotropy group $e^{ip}K^{\mathbb{C}}e^{-ip}$ or, equivalently, that for $u \in \mathfrak{p}^{\mathbb{C}}, (\text{ad } e^{-ip})u \notin \mathfrak{k}^{\mathbb{C}}$. We have

$$(\text{ad } e^{-ip})u = e^{\text{ad}(-ip)}u = \cosh(\text{ad}(-ip))u + \sinh(\text{ad}(-ip))u, \quad (3.4)$$

where the first term of the sum lies in $\mathfrak{p}^{\mathbb{C}}$ and the second one in $\mathfrak{k}^{\mathbb{C}}$. To show that the first term does not vanish recall that $(\text{ad}(-ip))^2$ has all eigenvalues nonnegative. $\square$

We now have:

Lemma 3.2. *In the complex coordinates $z = a + ib$ given by the previous lemma, the Levy form (3.3) of a G -invariant function w satisfies the equation:*

$$\left(\frac{\partial^2 w}{\partial z_k \partial \bar{z}_l} \right)_{\substack{a=0 \\ b=0}} = \frac{1}{4} \frac{\partial^2}{\partial b_k \partial b_l} w(e^{ib} e^{ip})_{b=0}. \quad (3.5)$$

Proof. The polar decomposition of $G^\mathbb{C}$ implies that e^{a+ib} can be uniquely written as ge^{iy} , where $g \in G$ and $y \in \mathfrak{g}$. Any G -invariant function on $G^\mathbb{C}/K^\mathbb{C}$ in a neighbourhood of e^{ip} is a function of y only. On the other hand, as $e^{2iy} = (e^x e^{iy})^* (e^x e^{iy}) = e^{-a+ib} e^{a+ib}$, it follows from the Campbell-Hausdorff formula that $y = b + [b, a]/2 + \text{higher order terms}$. Hence the matrix of second derivatives in (3.3) at e^{ip} (i.e. at $a = 0, b = 0$) is the same as the matrix of second derivatives of

$$(a, b) \mapsto e^{(ib + \frac{i}{2}[b, a])} e^{ip} \quad (3.6)$$

at $a = 0, b = 0$. We shall now show that for a G -invariant function w on $G^\mathbb{C}/K^\mathbb{C}$, this matrix of second derivatives is equal to the right-hand side of (3.5).

The Campbell-Hausdorff formula implies that up to order 2 in a, b , we have $e^{(ib + \frac{i}{2}[b, a])} = e^{ib} e^{\frac{i}{2}[b, a]}$. Set $c = [b, a]/2$, which is a point in $\mathfrak{k}$. We are going to show that modulo terms of order 2 in c (hence of order 4 in a, b), $e^{ic} e^{ip}$ is equal to $e^\rho e^{ip} e^{iq}$, where $\rho \in \mathfrak{g}$ and $q \in \mathfrak{k}$ are both linearly dependent on c . We note that this proves the lemma, as

$$e^{ib} e^\rho e^{ip} e^{iq} = e^\rho e^{ib+O(3)} e^{ip} e^{iq} = e^\rho e^{ib+O(3)} e^{ip}$$

in $G^\mathbb{C}/K^\mathbb{C}$, where $O(3)$ denotes terms of order 3 and higher in a, b .

We find q from the equation $\cosh \text{ad}(ip)(q) = c$, which can be solved uniquely as $\cosh \text{ad}(ip)$ is symmetric and positive-definite on $\mathfrak{k} \subset \mathfrak{g}$. We then put $\rho = -i \sinh \text{ad}(ip)(q)$. We observe that $\rho \in \mathfrak{g}$ and $e^{\rho-ic} = e^{ip} e^{-iq} e^{-ip}$, thanks to (3.4). Moreover, modulo terms quadratic in c , $e^\rho = e^{ic} e^{\rho-ic}$ and, consequently:

$$e^\rho e^{ip} e^{iq} = e^{ic} e^{\rho-ic} e^{ip} e^{iq} = e^{ic} (e^{ip} e^{-iq} e^{-ip}) e^{ip} e^{iq} = e^{ic} e^{ip},$$

again modulo terms quadratic in c . This finishes the proof of the lemma. $\square$

According to this lemma, we have to compute $\frac{\partial^2}{\partial b_k \partial b_l} w(e^{ib} e^{ip})_{b=0}$. Now, since $e^{ib} e^{ip} \in G^*$, $e^{ib} e^{ip} = k e^{iz}$, where $z = z(b) \in \mathfrak{p}$ and $k \in K$. As w is G -invariant, $w(e^{ib} e^{ip}) = w(e^{iz})$ and therefore

$$\frac{\partial^2}{\partial b_k \partial b_l} w(e^{ib} e^{ip})_{b=0} = \frac{\partial^2}{\partial b_k \partial b_l} w(e^{iz(b)})_{b=0}.$$

Thus we compute the matrix of second derivatives of a function defined on $\exp(i\mathfrak{p})$ in the coordinates given by $b \mapsto e^{ib} e^{ip} \mapsto e^{iz(b)}$. These, however, are the geodesic coordinates at the point e^{ip} in the symmetric space $K \backslash G^*$ (being translations of geodesics at $[1]$), and hence the matrix of second derivatives in these coordinates is equal to the Riemannian Hessian (2.1) for the symmetric metric on $K \backslash G^*$.

We obtain

Theorem 3.3. *Let w be a smooth G -invariant function on $X = G^\mathbb{C}/K^\mathbb{C}$ and let $\bar{w}$ be its restriction to the fiber $S = \exp(i\mathfrak{p})$ of (3.2) over $[1]$. Let g denote the*

symmetric metric on $S \simeq K \backslash G^*$. Then w is (strictly) plurisubharmonic if and only if $\bar{w}$ is (strictly) g -convex. Moreover, the following equality holds:

$$\partial\bar{\partial} \log \det Lw = \partial\bar{\partial} \log \widehat{\mathbf{M}}_g(\bar{w}),$$

where $\hat{u} : X \rightarrow \mathbb{R}$ is a G -invariant function such that $\bar{\hat{u}}$ is a given K -invariant function u on S . $\square$

We are now ready to prove Theorem 1. Recall that $X = G^{\mathbb{C}}/K^{\mathbb{C}}$ is a Stein manifold and so if ρ is an exact $(1, 1)$ form on X , then $\rho = -i\partial\bar{\partial}h$ for some function h . If ρ is G -invariant, then we can assume that h is G -invariant. We can restrict h to the fiber S defined in the last theorem and thanks to Corollary 2.2 we can find a strictly g -convex K -invariant smooth solution $\bar{u}$ to the equation (2.2) with $f = e^h$, where the metric g is the symmetric metric on $S \simeq K \backslash G^*$. We can extend this solution via G -action to a G -invariant function u on X . Theorem 3.3 implies now that u is strictly plurisubharmonic and that the Ricci form of the Kähler metric with potential u is ρ .

Acknowledgement. This work has been supported by an Advanced Research Fellowship from the Engineering and Physical Sciences Research Council of Great Britain.

REFERENCES

- [1] H. Azad and J.-J. Loeb ‘Plurisubharmonic functions and Kählerian metrics on complexification of symmetric spaces’, *Indag. Math.* (N.S.) 3 (1992), 365–375.
- [2] R. Bielawski ‘Kähler metrics on $G^{\mathbb{C}}$ ’, *J. Reine Angew. Math.* 559 (2003), 123–136.
- [3] R. Bielawski ‘Entire invariant solutions to Monge-Ampère equations’, math.AP/0212205, to appear in *Proc. Amer. Math. Soc.*
- [4] O. Biquard and P. Gauduchon ‘Géométrie hyperkählérienne des espaces hermitiens symétriques complexifiés’, *Sémin. Théor. Spectr. Géom.* 16, Univ. Grenoble I, 127–173.
- [5] P. Heinzner and A. Huckleberry ‘Analytic Hilbert quotients’, in: *Several complex variables (Berkeley, CA, 1995–1996)*, 309–349, Math. Sci. Res. Inst. Publ., 37, Cambridge University Press, Cambridge, 1999.
- [6] S. Helgason *Differential geometry, Lie groups, and Symmetric spaces*, Academic Press, New York, 1978.
- [7] L. Lempert and R. Szöke ‘Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundle of Riemannian manifolds’, *Math. Ann.* 290 (1991), 689–712.
- [8] G.D. Mostow ‘On covariant fiberings of Klein spaces’ *Amer. J. Math.* 77 (1955), 247–278.
- [9] M.B. Stenzel ‘Ricci-flat metrics on the complexification of a compact rank one symmetric space’, *Manuscripta Math.* 80 (1993), 151–163.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, GLASGOW G12 8QW, UK
E-mail address: R.Bielawski@maths.gla.ac.uk